

Biased galaxy formation in the fields of high-redshift AGN

J. A. Stevens,¹ M. J. Page,² F. J. Carrera,³ R. J. Ivison,⁴ and Ian Smail⁵

¹*Centre for Astrophysics Research, University of Hertfordshire, College Lane, Herts AL10 9AB, UK*

²*Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK*

³*Instituto de Fisica de Cantabria (CSIC-UC), Avienida de los Castros 39005 Santander, Spain*

⁴*UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK*

⁵*Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE, UK*

Abstract. We discuss preliminary results from our programme to map the fields of high-redshift AGN. In the context of the hierarchical models such fields are predicted to contain an over-density of young, luminous galaxies destined to evolve into the core of a rich cluster by the present epoch. We have thus imaged from submillimetre to X-ray wavelengths the few-arcmin scale fields of a small sample of high-redshift QSOs. We find that submillimetre wavelength data from SCUBA show striking over-densities of luminous star-forming galaxies over scales of ~ 500 kpc. Whilst many of these galaxies are undetected even in deep near-IR imaging almost all of them are detected by *Spitzer* at 4.5, 8.0 and 24 μm , showing that they have extremely red colours. However, they are not detected in our *XMM-Newton* observations suggesting that any AGN must be highly obscured. Optical-through-mid-IR SEDs show the redshifted 1.6 μm bump from star-light giving preliminary evidence that the galaxies lie at the same redshift, and thus in the same structure, as the QSO although this finding must be confirmed with photometric and/or spectroscopic redshifts.

1. Introduction

The popular hierarchical model of galaxy formation predicts that elliptical galaxies found today in the cores of rich clusters formed at high redshifts and at rare high-density peaks of the dark matter distribution. Within these regions, gas rich proto-galaxies merge together rapidly and form stars at a high rate. The same reservoir of gas used to build the stellar mass can also fuel the growth of the supermassive black holes (SMBH) found dormant in the centre of the galaxies at low redshift (Kauffmann & Haehnelt 2000). Since massive star formation is known to be a dusty phenomenon, the light from such young galaxies may well be highly obscured rendering the population inaccessible to optical techniques. Dust enshrouded star formation is, however, a luminous phenomenon in the submillimetre (hereafter submm) through far-IR waveband where the star-light absorbed by dust grains is re-emitted.

Unfortunately, it is not currently feasible at submm/far-IR wavelengths to perform a survey of a randomly selected region of sky over an area sufficiently large to ensure that rare structures in the early universe are contained within it. However, one method of locating such structures is to target the fields of high-redshift AGN. Given their huge luminosities, such objects must already contain a SMBH, and therefore must represent some of the most massive objects in existence at their epochs. They should thus act as signposts to the rare high-density peaks that we wish to study and can target with existing technology at submm wavelengths (Ivison et al. 2000; Stevens et al. 2003). This paper presents preliminary results from such a programme. We concentrate on two QSO fields at $1 < z < 3$, presenting submm results obtained with SCUBA on the JCMT and follow-up imaging at near-IR (UKIRT) and mid-IR (*Spitzer*) wavelengths.

2. Over-densities of luminous, extremely red star-forming galaxies

The two targets discussed here are the X-ray absorbed QSOs RX J094144.51 + 385434.8 ($z = 1.82$) and RX J121803.82 + 470854.6 ($z = 1.74$). The importance of these objects has been discussed extensively in the literature (Page et al. 2004; Stevens et al. 2005). See also the contribution by Page et al. in this volume.




Figure 1. Submm, mid- and near-IR imaging of the RX J121803.82 + 470854.6 field. Left-hand panel shows 850 μm contours (2,3,4,5 σ) on a deep K -band image whilst the right-hand panel shows the same contours on an 8.0 μm image. Note that dust peaks lacking K -band counterparts often do have detections at 8.0 μm . Each panel is ~ 1 arcmin square.

Submm, near- and mid-IR images of the RX J121803.82 + 470854.6 field are shown in Fig. 1. The 850 μm data reveal a large over-density of sources, particularly pronounced in the close vicinity (~ 30 arcsec) of the QSO. Results from blank-field surveys predict, at most, one $\geq 3 \sigma$ source in addition to the QSO. The near- and mid-IR imaging gives an indication that many of these submm sources are indeed real and have extremely red colours. Many of them are detected by *Spitzer* even when they are not seen in deep ($K \sim 20.5$) near-IR imaging; e.g. the two most significant dust peaks in Fig. 1. Star-formation

rates computed in the standard manner are at least several $100 M_{\odot} \text{ yr}^{-1}$. It can thus be concluded that we have discovered a large over-density of star-forming galaxies in the few hundred kpc-scale environment of the QSO.




Figure 2. Submm and mid-IR imaging of the RX J094144.51 + 385434.8 field. $450 \mu\text{m}$ contours on the $850 \mu\text{m}$ image (left), $450 \mu\text{m}$ contours on the $4.5 \mu\text{m}$ image (centre) and $450 \mu\text{m}$ contours on the $24 \mu\text{m}$ image (right). Submm contours start at 2σ and increase in 1σ steps. Panels are $\sim 1.0 \times 1.5$ arcmin. The structure of galaxies is ~ 400 kpc in extent (Stevens et al. 2004).

Fig. 2 shows submm and mid-IR data for the RX J094144.51 + 385434.8 field. Here we see a similar over-density of luminous star-forming galaxies to that discussed above. For this dataset the observing conditions were sufficiently transparent that we were able to collect good data at $450 \mu\text{m}$. These data show that the large-scale filamentary structure seen at $850 \mu\text{m}$ breaks up into individual star-forming galaxies when observed at higher resolution (~ 8 cf. ~ 14 arcsec). Such datasets were very rarely obtained with SCUBA at $450 \mu\text{m}$ because of the difficulty of observing at short submm wavelengths from Mauna Kea. However, the increased sensitivity and field-of-view afforded by SCUBA-2 will go a long way to alleviating this problem and we can look forwards to higher resolution images of distant galaxies.

A second result is that the submm-selected galaxies are detected by *Spitzer* with both IRAC and MIPS (Fig. 2) verifying their reality and, with the addition of optical and near-infrared imaging, allowing us to construct SEDs of the individual galaxies (see Fig. 3). We find that the SEDs are quite similar to that of the local ULIRG Arp 220 if it is moved to the redshift of the QSO ($z = 1.82$) suggesting that the bump we see in the spectra, peaking roughly in the $4.5 \mu\text{m}$ band is the redshifted $1.6 \mu\text{m}$ bump from star-light. This result provides the first evidence that the luminous star-forming galaxies discovered by SCUBA are at the same redshift and in the same structure as the QSO. We are currently working to confirm this finding with photometric and spectroscopic redshifts.

2.1. Buried AGN?

If these galaxies are to evolve into cluster elliptical galaxies then the SMBHs found at their centres at the current epoch should be active at $z \sim 2$. Our 50 ks *XMM-Newton* observations detect the QSOs but not the star-forming galaxies in




Figure 3. Optical through mid-IR spectra of the QSO RX J094144.51 + 385434.8 (top dotted line and points) and the submm galaxies in its vicinity. The solid line is the spectrum of the local ULIRG Arp 220 redshifted to $z = 1.82$ and scaled arbitrarily (Silva et al. 1998).

their fields. If we assume that these galaxies contain an AGN with a luminosity $L(2 - 10 \text{ keV}) = 10^{44} \text{ erg s}^{-1}$ and a photon index $\Gamma = 2$ then the non-detections imply a column density $N_H > 5 \times 10^{23} \text{ cm}^{-2}$. Therefore, if these galaxies contain reasonably powerful AGN they must be highly obscured.

3. Looking ahead to SCUBA-2 and *Herschel*

The next year will see major new facilities operating at far-IR and submm wavelengths. Both *Herschel* and SCUBA-2 will provide much faster mapping speeds than those currently available allowing us to map an area of sky around the QSOs large enough to observe the whole filamentary structure of the collapsing proto-cluster rather than just the core region. Simulations show that this will require a map of at least 20 Mpc square at $z = 2$ (comoving units). At present a significant amount of telescope time is required to map one hundredth of this area but SCUBA-2 can map a region this large in only a few hours.

References

- Ivison, R. J., Dunlop, J. S., Smail, Ian, Dey, Arjun, Liu, M. C., & Graham, J. R., 2000, ApJ, 542, 27
- Kauffmann, G., & Haehnelt, M., 2000, MNRAS, 311, 576
- Page, M. J., Stevens, J. A., Ivison, R. J., & Carrera, F. J., 2004, ApJ, 611, L85
- Page, M. J., Carrera, F. J., Ebrero, J., Stevens, J. A., & Ivison R. J., this issue
- Silva, L., Granato, G. L., Bressan, A., & Danese, L., 1998, ApJ, 509, 103
- Stevens, J. A. et al. 2003, Nat, 425, 264
- Stevens, J. A., Page, M. J., Ivison, R. J., Smail, Ian, & Carrera, F. J., 2004, ApJ, 604, L17

Stevens, J. A., Page, M. J., Ivison R. J., Carrera, F. J., Mittaz, J. P. D., Smail, Ian, & McHardy, I. M., 2005, MNRAS, 360, 610